



## Journal of the Chemical Society, Perkin Transactions 2

[Issue 12, 1991](#)

Perkin Transactions 2 ceased publication in 2002. Organic & Biomolecular Chemistry is the RSC's organic chemistry journal for high-quality, original organic chemistry research.

### **EPR study of radical intermediates from the oxidation of 6-ethoxy-2,2,4-trimethyl- and 6-ethoxy-2,2,4,8-tetramethyl-1,2-dihydroquinoline**

[Frank D. Gunstone](#), [Raphael C. Mordi](#), [Snorri Thorisson](#), [John C. Walton](#) and [Richard A. Jackson](#)

*J. Chem. Soc., Perkin Trans. 2*, 1991, 1955-1958

DOI: 10.1039/P29910001955

[Share on citeulike](#) | [Share on facebook](#) | [Share on twitter](#) |

[More](#)

- PDF
- Article References
- [Add PDF to Basket \(£38\\*\)](#)
- \*Exclusive of taxes  
This article contains 4 page(s)

- [Abstract](#)

The EPR spectra of 1,2-dihydro-6-ethoxy-2,2,4-trimethyl- and 1,2-dihydro-6-ethoxy-2,2,4,8-tetramethyl-quinolin-1-yl radicals were observed in heptane solution. The hyperfine splittings showed that this class of radical is extensively delocalised with significant spin density at C(8). Both radicals decayed by second-order processes, the rate constants being  $5 \times 10^6$  and  $4 \times 10^2 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ , respectively at 273 K. The latter reaction is much slower because the 8-methyl substituent blocks the formation of the 1,8'-dimer. Both radicals reacted with oxygen to give the corresponding nitroxides, although reaction was very slow for the 8-methyl

derivative. A mechanism is proposed to rationalise product formation from 1,2-dihydro-6-ethoxy-2,2,4-trimethylquinoline when used as an antioxidant.

